## A Selection of SUNSHINE COAST BIRDS

The Living Smart program promotes the creation and protection of habitat for local native wildlife in urban backyards through its Backyard Biodiversity module. Wherever you live we can all play a part in helping our local native animals by providing safe, suitable habitat and feeding opportunities for them.





http://www.birdsaustralia.com.au

http://www.birdsinbackyards.net

nbackyards.net http://



http://www.sunshinecoast.qld.gov.au

your environment levy your habitat

Sunshine Coast

Council

Photographers: W Dabwroka and K Vang © Bird Explorers (1-3, 6, 10, 13, 14, 16, 18, 20, 26, 27, 29, 30, 33, 35, 36, 38, 41, 43, 45, 49, 52, 55, 56, 58–61, 63, 65, 66, 68, 71, 72, 76, 78); R Major © Australian Museum (7, 17, 19, 22, 23, 47, 48, 54, 57, 80); T Oliver (4, 9, 32, 37, 44, 53, 64, 67, 69, 75, 77); Á Lumnitzer (21, 34, 46, 51, 73, 79); E Pierce (28, 31, 62); N Caldwell (15, 42); M Barlow (5); J Irvine (8); A Beleski (11); A Los Baños (12); G Gowing © Australian Museum (24); A Kittilia (25); I and J Brown (39); B Gow (40); G Gates (50); B Ravich (70); C Tzaros (74)



**1. Pacific Black Duck** (55cm) Common duck often found on lawns and around lakes.



**2. Australian Wood Duck** (47 cm) Seen on grassland and away from water. Nests in hollows.



**3. Australian Brush-turkey** (65 cm) Large black ground forager. Males build mounds for females to lay eggs. Common in gardens.



**4. Brown Quail** (18 cm) Largest native quail. Found in groups in grasses near water.



5. White-faced Heron (65 cm) Most common heron. Can be found anywhere with water. Often seen on lawns.



**6. Great Egret** (80 cm) Hunts alone in water but nests in a treetop colony.



**7. Australian White Ibis** (72 cm) Common in towns and cities. Will roost in palms and can become a pest.



8. White-bellied Sea Eagle (80 cm) Very large eagle that soars in large circles. 'Honking' call.



**9. Brahminy Kite** (50 cm) Found along coasts and rivers. Distinct white head.



**10. Whistling Kite** (55 cm) Shaggy appearance. Underwings have pale 'M' shape.



**11. Osprey** (66 cm) Large fishing hawk found on coastline. Bold brown eye stripe.



**12. Australian Hobby** (33 cm) Small, fast, dark falcon with square tail. Takes prey on wing.



**13. Purple Swamphen** (46 cm) Large conspicuous hen with bright red beak and legs.



**14. Dusky Moorhen** (37 cm) Medium sized dark hen with red beak. Swims but is unable to dive.



**15. Bush Stone-curlew** (55 cm) Unusual, bush dwelling wader. Wailing call at night. Skulks when alarmed.



**16. Masked Lapwing** (36 cm) Large, conspicuous, noisy and aggressive plover. Common around lawns and parks with short grass.



**17. Rock Dove\*** (34 cm) Introduced species also called the Feral Pigeon. Originally from Europe.



**18. Bar-shouldered Dove** (28 cm) Commonly seen feeding on the ground. Builds flimsy nests.



**19. Crested Pigeon** (32 cm) Grey pigeon with distinct black crest. Feeds on the ground. Distinct whistling sound on take off.



**20. Spotted Dove\*** (30 cm) Previously known as Spotted Turtledove. Introduced from Asia around 1860. Mostly in parks and gardens.



**21. Superb Fruit-dove** (23 cm) Small colourful dove. Rainforest and fringes, lantana thickets.



**22. Sulphur-crested Cockatoo** (48 cm) Familiar, noisy white cockatoo. Often feeds on ground. Can be destructive.



**23. Galah** (36 cm) Widespread and common bird. Feeds on ground or low foliage in small flocks.



**24. Yellow-tailed Black-Cockatoo** (60 cm) Feeds in small to large noisy flocks on native seeds, trees and pine cones.



**25. Glossy Black-Cockatoo** (48 cm) Classified as Vunerable. Reliant on casuarina seeds.



**26. Rainbow Lorikeet** (30 cm) Familiar colourful lorikeet. Feeds on fruit and nectar from flowers.



**27. Scaly-breasted Lorikeet** (23 cm) Feed in flocks, sometimes with rainbow lorikeets. Found in well-treed suburbs.



**28. Pale-headed Rosella** (31 cm) Quiet parrot. Feeds mostly on seeds and sometimes fruits in grassy woodlands and trees near water.



**29.** Australian King-parrot (42 cm) Males are bright red (shown), females have a green head .



**30. Channel-billed Cuckoo** (62 cm) Huge, pale grey with large curved bill. Migratory species, spring/summer.



**31. Common Koel** (42 cm) Males are black (shown), females and young are mottled brown. Migratory species spring/summer.



**32.** Pheasant Coucal (60 cm) Carnivorous species with long tail. The only Australia cuckoo to build its own nest.



**33. Fan-tailed Cuckoo** (26 cm) Hunts from perch, dropping on prey. Has a mournful descending call.



**34. Southern Boobook** (29 cm) Also called a 'Mopoke'. Smallest and most common owl in Australia.



**35. Tawny Frogmouth** (44 cm) Nocturnal bird that feeds on invertebrates and some small animals. Usually seen in camouflage pose on bare branch.



**36. Laughing Kookaburra** (42 cm) Nests in hollows and termite nests. Uses its large beak to catch reptiles and small snakes. Classic laughing call.



**37. Sacred Kingfisher** (22 cm) Hunts on dry land for small reptiles. Margins of rivers and open woodland.



**38. Rainbow Bee-eater** (24 cm) Orange wings flash against the sky. Tail has long central streamers.



**39. Dollarbird** (28 cm) Arrives each year in September to breed. Uses hollows for nesting.



**40. Red-backed Fairy-wren** (11 cm) Females and young are brown. Found in tall grass and shrubs.



**41. Variegated Fairy-wren** (13 cm) Females and young are brown. Shy, inclined to stay in undergrowth.



42. Blue-faced Honeyeater (31 cm) Large, gregarious, aggressive. Adults have a blue mask, young have a green mask.



**43. White-cheeked Honeyeater** (18 cm) White cheeks and heavy dark bill. Often in groups foraging for nectar.



**44. Lewin's Honeyeater** (20 cm) Yellow ear patch. Feeds on fruit, nectar and on insects in trees.



**45. Brown Honeyeater** (14 cm) Brown with small yellow tuft behind eye. Feeds on flowering natives.



**46. Scarlet Honeyeater** (10 cm) Females and young are brown. Hovers at blossoms to feed.



**47. Noisy Miner** (25 cm) Aggressive honeyeater that lives in large family groups. Territorial.



**48.** Noisy Friarbird (34 cm) Conspicuous, black head without feathers, has bump on bill. Noisy abrupt call when feeding.



**49. Striated Pardalote** (9 cm) Feed on insects in the tree canopy. Nest in hollows or tunnel in earthy banks.



**50. Brown Thornbill** (10 cm) Warm brown to olive brown in colour. Actively seeks insect prey in undergrowth.



**51. Striated Thornbill** (10 cm) Small insect-eater with an orangebrown cap. Feeds high up in trees.



**52. Eastern Whipbird** (28 cm) Loud whip-crack call made by 2 birds. Keeps to thickets, fossicking in ground litter.



53. Torresian Crow (53 cm) Forms large noisy groups in urban areas. Nest of sticks high in eucalypt.



**54. Pied Currawong** (46 cm) Bold scavengers in suburbs. Form large, often noisy, wandering flocks.



**55. Grey Butcherbird** (27 cm) Melodic song. Aggressive predator that stores food on tree branches or crevices.



**56. Pied Butcherbird** (36 cm) Black-headed with superb song. Hunts from bare limbs, power poles and wires.



57. Australian Magpie (44 cm) One of Australia's best known birds. Conspicuous, large, black and white,



58. Golden Whistler (18 cm) Persistent, loud, ringing song, Colourful male, solitary except when breeding.



59. Rufous Whistler (18 cm) Back, tail and wings mostly midgrey, with deep rusty yellow/orange underparts unlike the bright yellow of the Golden Whistler.



60. Grey Shrike-thrush (25 cm) Rich, varied song, Widespread in extremely varied habitats.



61. Black-faced Cuckoo-shrike (36 cm) Conspicuous black face and throat. Strong undulating flight aids identification.



62. Australasian Figbird (29 cm) Lives in small groups. Feeds on fruiting trees, wild figs and exotic palm fruits.



63. Olive-backed Oriole (28 cm) Graceful thrush-like bird with greenish and streaked plumage. Difficult to sight.



64. Spangled Drongo (32 cm) Iridescent black plumage, with long, out curved, forked tail. Upright posture.



65. Willie Wagtail (22 cm) Waves tail incessantly. Always restless, swinging body from side to side while making a clicking sound.



66. Grey Fantail (17 cm) Grey with short white eyebrow. Twisting, turning acrobatics when feeding.



67. Rufous Fantail (16 cm) Flits and dances above lower undergrowth. Fanned tail bright orangerufous.



68. Magpie-lark (30 cm) Common magpie-like small bird. Has adapted to use man-made environments.



69. Leaden Flycatcher (16 cm) Hovers above foliage, taking small insects from leaves or snatching in flight.



70. Eastern Yellow Robin (16 cm) Clings to low branches and tree trunks while scanning the ground for movement of insects.



71. White-breasted Woodswallow (18 cm) Grev hood and white breast. Found in woodlands near water or mangroves. Highly sociable.



72. Welcome Swallow (15 cm) Slender, streamlined with long tapered wings. Fast, buovant, acrobatic in air.



73. Fairy Martin (13 cm) Catches flying insects in flight. Builds bottle-shaped mud nest under any overhang.



74. Silvereye (13 cm) Lively, busily active little birds, constantly on the move. Feed on fruit and insects.



75. Red-browed Finch (12 cm) Highly sociable, with scarlet bill, highbrow and rump. Perches on grass.



76. Double-barred Finch (11 cm) Feeds on grass seeds and flits about in low shrubs.



77. Chestnut-breasted Mannikin (12 cm) Strong silver bill/black face with chestnut upper breast. Seed eater.



**78. Common Myna\*** (25 cm) Also called the Indian Myna. Introduced to N Qld in late 1800's and classified as a pest.



79. Common Starling\* (21 cm) Introduced to eastern states in late 1800's. Flight swift, flocks rise and fall.



80. House Sparrow\* (16 cm) Introduced in the 1860's, now an abundant pest.

